

The actions, detection and synthesis of platelet-activating factor (PAF) in mammalian pregnancy

Liu Yutang (刘玉堂)

College of Wildlife Resources, Northeast Forestry University, Harbin 150040, P. R. China

Sui Shuqin (隋淑芹)

(Harbin Grain Bureau)

Sun Zhongwu(孙中武) Bi Bing (毕冰) Xu Cuiqing(许翠清)

College of Wildlife Resources, Northeast Forestry University, Harbin 150040, P. R. China

Abstract Platelet-activating factor (PAF) exhibits a variety of biological activities and it be thought to involved in various pathophysiological process. In this paper, some studies were summarized about those roles of PAF in a variety productive processes of female of mammalian that include fertilization, implantation and parturition, and that was involved in the concentration, synthesis, degradation and some assay methods of PAF. The relationship between PAF and early pregnancy factor (EPF) was reviewed.

Key words: Platelet-activating factor, Mammalian, Pregnancy, Early pregnancy factor

Platelet-activating factor (PAF:1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine) was discovered as a chemical mediator, released from sensitized basophils (Benveniste *et al* 1972), that caused platelet aggregation. Subsequent studies demonstrated that PAF was produced upon appropriate stimulation of a variety of human cells, including neutrophils, eosinophils, monocytes, and endothelial cells, and this potent lipid mediator is involved in a number of the allergic and inflammatory responses (Hanahan 1986; Braquet *et al* 1987; Wallace 1989; Prescott *et al* 1990). PAF is also found in spermatozoa (Kumar *et al* 1988) preimplanted embryos (O'Neill 1985), amniotic fluid (Billah 1983), and the uterus (Yasuda *et al* 1986; Angle *et al.* 1988). PAF is considered to be involved in a variety of reproductive processes including fertilization (Roudebush *et al* 1990), implantation (Ryan *et al* 1990), and parturition (Johnston 1989). The roles of PAF in parturition are not fully defined, but it is well documented that PAF induces myometrial contraction (Tetta *et al* 1986) and promotes the synthesis of prostaglandin E₂ (Billah *et al* 1985).

PAF in the establishment of pregnancy and ovoimplantation

PAF may be the first physiology signal produced by the embryo for maternal recognition of pregnancy. In mice and humans fertilization induces a pregnancy-associated thrombocytopenia (O'Neill 1985a) which has been attributed to release of platelet-activating factor (PAF) by the zygote (O'Neill 1985b, c). Ammit and O'Neill (1991) subsequently confirmed the release of embryonic PAF into culture media, and

O'Neill and Saunders (1984) identified a correlation between media PAF concentration, embryo implantation rate and successful pregnancy.

PAF is produced by uterine tissue and pre-implantation-stage embryos. PAF level in the rabbit uterus increase from Day 3 to Day 5 of pregnancy (Angle *et al* 1988). Embryonic PAF production in the rabbit and mouse also increases during the pre-implantation phase, with maximum levels at the expanded blastocyst stage (Minhas *et al* 1993; Ripps *et al* 1993). PAF production by human embryos has been correlated with pregnancy potential (Nakatsuka *et al.*, 1992; O'Neill *et al* 1987). Furthermore, mouse embryos cultured in the presence of PAF have enhanced developmental rates (Roberts *et al* 1993) and higher implantation rates upon transfer to synchronized recipients (Ryan *et al* 1987). Addition of exogenous PAF to the culture medium improves two-cell development to the blastocyst stage in the mouse (Nishi *et al* 1995; Roudebush *et al* 1996; O'Neill 1997), apparently which may be due, in part, to embryo-derived PAF stimulation of embryonic metabolism (Ryan *et al* 1989). PAF directly influences the oxidative metabolism of glucose and lactate in the pre-implantation mouse embryo (Ryan *et al* 1990). These observations suggest that PAF production is a prerequisite for pregnancy, and that it could be used as a marker for embryo viability. The effect of PAF on in vitro mouse embryo development appears to be strain specific (Radonjic-Lazovic *et al* 1995). Enhanced embryo development has also been reported in rabbit oocytes fertilized in vitro with PAF-treated spermatozoa (Roudebush *et al* 1993).

PAF antibodies inhibit embryo development (Roudebush *et al* 1993), and antagonists inhibit im-

plantation (Spinks *et al* 1988; Andu *et al* 1990; Norris *et al* 1994). Providing further evidence on the presence and requirement of embryo-derived PAF during the pre-implantation period. It appears that PAF's action in the embryo may be receptor mediated, since various PAF antagonists competitively inhibit its action (Nishi *et al* 1995). Roudebush *et al* (1997) determine the presence of the PAF receptor in the mouse preimplantation-stage embryo, and exogenous PAF treatment of the two-cell-stage CFW mouse embryo resulted in a fourfold increase in intracellular calcium over background levels, calcium has proven to be an important second messenger that controls or influences a variety of cellular functions. PAF receptors have been described in both sites in other species (Yang *et al* 1991, 1992).

Velasquez *et al* (1995) found that the PAF antagonists TCV-309 and BN-52021 delayed significantly the transport of egg to the uterus in pregnant animals, but not in cycling animals; i.e., they retarded the passage of embryos but not of oocytes to the uterus. Administration of PAF to cycling hamsters hastened the oviductal transport of ova. These data suggest that in the hamster, the earlier passage of embryos to the uterus as compared to oocytes is mediated by PAF. The decidual-like reaction induced by PAF is observed by Acker *et al* (1989) in the rat uterus.

Taken together those all confirmed that PAF is an important mediator in the establishment of pregnancy and ovo-implantation which may lead to improvement in *in vitro* fertilization and open up the possibility that PAF antagonists may be used as local contraceptive agents.

The Relationship between PAF and EPF

Early pregnancy factor (EPF) is one of the earliest biochemical indicator of pregnancy (Morton *et al* 1974). It has been detected at about the time of the pronuclear stage of the embryo in all mammals investigated including mice (Morton *et al* 1974), human (Morton *et al* 1977), several domestic (Morton *et al* 1979; Nancarrow *et al* 1981) and other species. The ability of EPF in sera of pregnant animals to act on lymphocytes to cause increased rosette inhibition titres in the rosette inhibition assay has been studied for a considerable time (Smart *et al* 1981; Whyte and Heap 1983; Morton 1984; Chard and Grudzinkas 1987; Morton *et al* 1987).

Orozco *et al* (1986) injected PAF into mature female mice at fourth stage of the oestrous cycle, mature male mice and immature female mice. Only animals that were in dioestrus, pro-oestrus or oestrus gave a positive EPF result within 1 h of injection. This group also demonstrated that PAF in combination with oestrous mouse serum could give a positive

response in the rosette inhibition test (Orozco *et al* 1990). The susceptibility of EPF activity in sera of pregnant animals to ammonium sulfate fractionation is well known (Clarke *et al* 1980; Sueoka *et al* 1989). Clark *et al* (1994) reinvestigated the effects of ammonium sulfate fractionation. Treatment of sera from pregnant mice with 40% ammonium sulfate was shown to liberate low molecular mass active moieties from their association with macromolecular serum components. After centrifugation, these active moieties partition into the supernatant fraction, while the macromolecular components to which they are bound partition into the pellet fraction. The macromolecular components of the supernatant and pellet fraction freed of any association with these low molecular mass moieties by extensive dialysis, can not alone induce increased rosette inhibition titres. In combination, however, components in the dialyzed pellet retentate fraction, cooperate with components in the dialysed supernatant retentate fraction to allow the expression of increased rosette inhibition titres when applied to fresh spleen cell in the rosette inhibition assay. In two-step incubation protocols, a prescribed order of addition must be followed if this cooperative effect is to be observed, namely supernatant retentate fraction in the second step, but not vice versa (Clark *et al* 1994).

Orozco *et al* (1994) demonstrated that the supernatant retentate fractions derived from sera of pregnant mice were functionally equivalent to PAF, or a serum stimulus because they stimulated the spleen cells used in the assay to produce active moieties and cooperated with pure thioredoxin in allowing for expression of activity. Conversely, the pellet retentate fractions obtained from sera of pregnant mice were shown to be functionally equivalent to thioredoxin in that they cooperated with a PAF stimulus to allow for the expression of increased rosette inhibition titres. Lash *et al* (1997) confirmed that PAF could stimulate the production of EPF by oestrous mouse ovaries and oviducts *in vitro*. Taking together the results that it could be suggested that PAF may be a component part responsible for EPF activity or PAF has the ability to induce the synthesis of EPF.

Biosynthesis and Degradation of PAF

PAF concentration in tissues or plasma is governed by the equilibrium between its biosynthesis and degradation. Studies in various cell types indicate that PAF synthesis may occur via one of two biosynthetic pathways the 'de novo' and 'membrane remodeling' pathways (for review see Snyder 1990). The immediate precursors of PAF in the de novo pathways are 1-O-alkyl-2-acetyl-sn-glycerol (alkylacetyl-glycerol) and cytidinediphosphocholine. In the membrane re-

modeling pathway, the substrates are 1-O-alkyl-2-lyso-sn-glycero-3-phosphocholine (lyso PAF) and acetyl-coenzyme A (acetyl-CoA). These reactions are catalyzed by dithiothreitol-insensitive cytidinediphosphocholine: 1-O-alkyl-2-acetyl-sn-glycerol cholinephosphotransferase (EC 2.7.8.16, cholinephosphotransferase), and by acetyl-coenzyme A:1-O-alkyl-2-lyso-sn-glycero-3-phosphocholine acetyltransferase (EC 2.3.1.67, acetyltransferase), respectively (Snyder 1990). The net synthesis of PAF occurs via the de novo pathway. The remodeling pathway, on the other hand, plays a role in the acute inflammatory process (Wykle *et al* 1980; Renooij *et al* 1981). Preimplantation embryos of mice (O'Neill 1985), humans (Collier *et al* 1988, 1990) and sheep (Battye *et al* 1991) secrete platelet-activating factor, which is detected in culture media. Production of PAF by the embryo in culture media ranges from 4.8 to 385.8 ng PAF per-embryo per 24 h for human embryos created by in vitro fertilization, and from 0.8 to 16.9 ng PAF per embryos per 24 h for mouse embryos (Collier *et al* 1988). This high output suggests that pre-implantation embryos may produce PAF by synthesis de novo.

Wells and O'Neill (1994) successfully detected the PAF biosynthetic enzymes, cholinephosphotransferase and acetyl-transferase in extracts of oocytes, zygotes and per-implantation embryos. The results indicate that the enzymes that catalyze the final step in both biosynthetic pathway are present within the oocyte and early embryo, while the activation of acetyltransferase following fertilization indicates that this enzyme may be of regulatory significance in the initiation of biosynthesis of PAF by zygotes.

PAF is generally thought to be labile in vivo. Biologically active PAF is converted to the biologically inactive lyso-PAF by PAF-acetylhydrolase (PAF-AH). PAF-AH activity is decreased in maternal plasma in rabbits (Maki *et al* 1988), human (Johnston 1989), rat and in rat uterus, in contrast, the placental PAF-AH activity significantly increased during the later stages of pregnancy (Matsubara *et al* 1997). It has been suggested that PAF-AH may regulate, in part, the PAF concentration in plasma or tissue.

The Measure of PAF

PAF bioassay

Embryo-derived PAF was discovered by O'Neill (1985), using a splenectomized mouse bioassay. Briefly, medium or test substances were given as a single i. p. injection (300 μ l) into splenectomized mice. Platelet counts were performed immediately before and 30 min after injection, animals were bled under light anaesthesia from the peri-orbital plexus and 10 μ l blood were diluted into 2 ml ammonium

oxalate (1% w/v). In each assay, samples of PAF-acether were run as positive controls.

Collier *et al* (1988) provides a quantitative rapid (15 min) method to measure PAF based on the determination of PAF-induced platelet aggregation. A platelet analyzer was used to count single platelet in a suspension of whole blood.

The another traditional methods for the detection and measurement of PAF rely on the release of tritiated serotonin (Hanahan and Weintraub 1985)

PAF radioimmuno assay

Smal believed that all those biological assays have problems of large variability within and between assays, and the need for careful control of non-specific interference which cause platelet activation. The development of a specific radioimmunoassay (RIA) for PAF (Smal *et al* 1990b, c) overcomes some of these problems, and allows accurate quantitation of low levels of PAF in biological samples (0.02-2 pmol). Briefly, the acidified sheep anti-PAF antiserum, the anti-sheep Ig G, synthetic PAF standard or unknown acidified sample and the (125 I)-labelled PAF tracer were mixed together. The radioactivity was quantitated by RIA. The PAF concentration was counted.

References

- Acker, G., Braquet, P. and Mencia-Huerta, J.M. 1989. Role of platelet-activating factor (PAF) in the initiation of the decidual reaction in the rat. *J Reprod Fertil* **85**:623-629
- Ammit, A. J. and O'Neill, C. 1991. Comparison of a radioimmunoassay and bioassay for embryo-derived platelet-activating factor. *Hum Reprod*, **6**: 872-878
- Andu, M., Suginami, H., and Matsuura, S. 1990. Pregnancy suppression by a structurally related antagonist for platelet-activating factor. *Asia Oceania. J Obstet Gynecol (Jpn)* **16**: 283-290
- Angle, M.J., Jones, M.A., McManus, L.M., Pinckard, R.N. and Harper, M.F.K. 1988. Platelet-activating factor in the rabbit uterus during early pregnancy. *J Reprod Fertil*, **83**: 711-722
- Battye, K. M., Ammit, A.J., Evans, G. and O'Neill, C. 1991. Production of platelet activating factor in vitro by preimplantation sheep embryos. *J Reprod Fertil*, **93**: 507-514
- Benvenist, J., Henson, P.M. and Cochrane, C.G. 1972. Leukocyte-dependent histamine release from rabbit platelets. The role of IgE, basophils and platelet-activating factor. *J Exp Med* **136**:1356-1377
- Billah, M. M., DiRenzo, G.C., Ban, C., Ruong, C.T., Hoffman, D.R., Anceschi, M.M., Bleasdale, J.E. and Johnston, J.M. 1985. Platelet-activating factor metabolism in human amnion and the responses of this tissue to extra cellular platelet-activating factor. *Prostaglandins*, **30**: 841-850
- Billah, M. M. and Johnston, J. M. 1983. Identification of

- phospholipid platelet-activating factor (1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholin) in human amniotic fluid and urine. *Biochem Biophys Res Commun*, **113**:51-58
- Braquet, P., Touqui, L., Shen, T. Y. Vargaftig, B.B. 1987. Perspectives in platelet-activating factor research. *Pharmacol Rev*, **39**: 97-145
- Chard, T. and Grudzinkas, J.G. 1987. Early pregnancy factor. *Biol Res Progress*, **8**: 53-56
- Clake, F.M, Morton, H., Rolfe, B.E. and Clunie, G.J.A. 1980. Partial characterisation of early pregnancy factor in sheep. *J Reprod Immun*, **2**: 151-162.
- Clark, I.Q, Orozco, C, Cock, I.E. and Clarke, F.M. 1994. The 'early pregnancy factor' revisited: the effect of ammonium sulfate on the capacity of pregnant mouse sera to express activity in the rosette inhibition assay. *J Reprod Fertil*, **100**: 279-289.
- Collier, M., O'Neill, C., Ammit, A.J. and Saunders, D. M. 1988. Biochemical and Pharmacological characterisation of human embryo-derived platelet activating factor. *Hum Reprod*, **3**: 993-998.
- Collier, M., O'Neill, C., Ammit, A. J. and Saunders, D. M. 1990. Measurement of human embryo-derived platelet-activating (PAF) using a quantitative bioassay of platelet aggregation. *Hum Reprod*, **5**: 323-328
- Hanahan, D. J. and Wientraub, S.T. 1985. Platelet activating factor; isolation, identification and assay. *Methods Biochem Anal*, **31**:195
- Hanahan, D. J. 1986. Platelet-activating factor: a biologically active phosphoglyceride. *Annu Rev Biochem* **55**: 483-509
- Johnston, J.M. 1989. The function of PAF in the communication between the fetal and maternal compartments during parturition. In: Saito K. Hanahan DJ (eds.), *Platelet-Activating Factor and Diseases*. Tokyo: International Medical Publishers, 129-138
- Kumar, R., Harper, M.J.K and Hanahan, D. J. 1988. Occurrence of platelet-activating factor in rabbit spermatozoa. *Arch Biochem Biophys*, **260**: 497-502.
- Lash, G.E and Legge, M. 1997. Induction of early pregnancy factor activity in vitro by platelet-activating factor in mice. *J Reprod Fertil*, **109**: 187-191
- Maki, N., Hoffman, D.R., and Johnston, J.M. 1988. Platelet-activating factor acetylhydrolase activity in maternal, fetal, and newborn rabbit plasma during pregnancy and lactation. *Proc Natl Acad Sci USA*, **85**: 728-732
- Matsubare, T., Yasuda, K., John, J.M., Sanesumi, M., Okada, H., Matsuoka, S., and Kanzaki, H. 1997. Platelet-activation factor (PAF) and PAF acetylhydrolase activity in rat uterus and placenta during the late stages of pregnancy. *Biol Reprod*, **56**: 885-890
- Minhas, B.S., Zhu, Y.P., Kim, H.N., Burwinkel, T.H, Ripps B.A., Buster, J.E. 1993. Embryonic platelet activating factor production in the rabbit increase during the pre-implantation phase. *J Assist Reprod Genet*, **10**: 366-370
- Morton, H, Hegh, V. and Clunie, G.J.A. 1974. Immunosuppression detected in pregnant mice by rosette inhibition test. *Nature*, **249**: 459-460
- Morton, H., Rolfe, B., Clunie, G.J.A., Anderson, M.J. and Morrison, J. 1977. An early pregnancy factor detected in human serum by the rosette inhibition test. *Lancet*, **1** 304-397
- Morton, H., Nancarrow, C. D., Saramuzzi, R.J, Evison, B.M and Clunie, G.J.A. 1979. Detection of early pregnancy in sheep by the rosette inhibition test. *J Reprod Fertil*, **56**: 75-80
- Morton, H. 1984. Early pregnancy factor (EPF): a link between fertilization and immunomodulation. *Austr J Biol Sci* **37**: 393-407
- Morton, H., Rolfe, B. and Cavanagh, A. C. 1987. Ovum factor and early pregnancy factor, *Current Topics in Dev Biol*, **23**: 73-92
- Nancarrow, C.D, Wallace, A.L.C and Grewal, A. S. 1981. The early pregnancy factor of sheep and cattle. *J Reprod Fertil (Suppl)*, **30**: 191-199
- Nakatasuka, M., Yoshida, N., Kudo, T. 1992. Platelet activating factor in culture media as an indicator of human embryonic development after in vitro fertilization. *Hum reprod* **7**: 1435-1439
- Nishi, O., Tominaga, T. , Gotor, Y. , Hayashi, K., and Mori, T. 1995. Effects of platelet activating factor on mouse embryo implantation in vitro. *J Assist Reprod Genet* **12**: 330-334
- Norris, C.J, Peairs, W.A., Kudolo, G.B, Newton, E. R. and Harper, M.J.K. 1994. Platelet-activating factor antagonists and implantation in rabbits. *J Reprod Fertil*, **100**: 395-401
- O'Eill, C. 1985a. Partial characterization of the embryo-derived platelet-activating factor in mice. *J Reprod Fertil* **75**: 375-380
- O'Neill, C. 1985b. Thrombocytopenia is an initial maternal response to fertilization in mice. *J Reprod Fertil (Suppl)* **30** 191-199
- O'Neill, C. 1985c. Examination of the cause of early pregnancy-associated thrombocytopenia in mice. *J Reprod Fertil*, **73**: 559-566
- O'Neill, C. 1995. Platelet-activating factor-antagonists reduce implantation in mice at low doses only. *Reprod Fertil Dev*, **7**: 51-57
- O'Neill, C. and Saunders, D. M. 1984. Assessment of embryo quality. *Lancet*, **2** 1035
- O'Neill, C., Gidley-Baird, A. A., Ammit, A.J., and Saunders, D.M. 1987. Use of a bioassay for embryo-derived platelet-activation factor as a means of assessing quality and pregnancy potential of human embryo. *Fertil Steril*, **47**:967-975
- O'Neill, C. 1997. Evidence for the requirement of autocrine growth factors for development of mouse preimplantation embryos in vitro. *Biol Reprod*, **56**:229-237
- Orozco, C., Perkins, T. and Clarke, F.M. 1986. Platelet-activating factor induces the expression of early pregnancy factor activity in female mice. *J Reprod Fertil*, **78**: 549-555
- Orozco, C., Cock, I., Perkins, A.V. and Clake, F. M. 1990. Platelet-activating factor and serum components from oestrous mice co-operate to mimic the activity of 'early pregnancy factor' in the rosette inhibition assay. *J Re-*

- prod Fertil, **88**: 447-457
- Orozco, C., Clark, I.Q., Cock, I.E. and Clarke, F.M. 1994. A description of the basic system of components in pregnant mice sera responsible for early pregnancy factor activity. *J Reprod Fertil*, **100**: 291-297.
- Prescott, S.M., Zimmerman, G. A., McIntyre, T. M. 1990. Platelet-activating factor. *J Biol Chem*, **265**: 17381-17384.
- Radonjic-Lazovic G, Roudebush, W. E. 1995. The effect of short-vs. long term platelet activating factor exposure on mouse pre-implantation embryo development. Early pregnancy. *Biol Med*, **1**:196-200
- Renooij, W. and Snyder, F. 1981. Biosynthesis of 1-alkyl-2-acetyl-sn-glycerol-3-phosphocholine (platelet activating factor and a hypotensive lipid) by cholinephosphotransferase in various rat tissues. *Biochim Biophys Acta*, **663**:545-556
- Ripps, B. A, Zhu, Y. P, Burwinkel, T. H., Kim, H.N., Buster J.E. and Minhas, B. S. 1993. Platelet-activation factor production from vitro and in vitro fertilized murine embryos is similar. *Am J Reprod Immunol*, **30**: 101-104
- Roberts, C., O'Neill, C., Wright, L. 1993. Platelet-activating factor(PAF) enhances mitosis in preimplantation mouse embryos. *Reprod Fertil Dev*, **5**: 271-279
- Roudebush, W.E., Minhas, B.S., Ricker, D.D., Palmer, T.V., Dodson, M.G. and Fortunato, S.J. 1990. Platelet-activating factor enhances in vitro fertilization of rabbit oocytes. *Am J Obstet Gynecol*, **163**:1670-1673
- Roudebush, W.E., Fukuda, A. I. and Minhas, B. S. 1993. Enhanced embryo development of rabbit oocytes fertilized in vitro with platelet-activating factor treated spermatozoa. *J Assist Reprod Genet*, **10**: 91-94
- Roudebush, W.E., Duralia, D.R and Butler, W.J. 1996. Effect of platelet activating factor(PAF) on preimplantation mouse B₆D₂F₁/J embryos formation. *Am J Reprod Immunol*, **35**:272-276.
- Roudebush, W. E., LaMarche, M.D, Levine, A.S, Jiang, H.X, and Butler, W.J. 1997. Evidence for the presence of the platelet-activating factor receptor in the CFW mouse preimplantation two-cell-stage embryo. *Biol Reprod*, **57**:575-579.
- Ryan, J. P, Spinks, N. R., O'Neill, C., Saunders, D.M. and Wales, R.G. 1987. Enhanced rates of implantation and increased metabolism of mouse embryos in the presence of platelet-activating factor. *Proc Fertil Soc Austr*, **6**:47
- Ryan, J. P, Spinks, N. R, O'Neill, C., Ammit, A.J and Wales R.G. 1989. Platelet-activating factor (PAF) production by mouse embryos in vitro and its effect on embryonic metabolism. *J Cell Biochem*, **40**: 387-395.
- Ryan, J. P., Spinks, N. R, O'Neill, C. and Wales, R.G. 1990. Implantation potential and fetal viability of mouse embryos cultured in media supplemented with platelet-activating factor. *J Reprod Fertil*, **89**:309-315.
- Smal, M.A., Baldo, B.A. and McCaskill, A. 1990. A specific, sensitive radioimmunoassay for platelet-activating factor (PAF). *J Immunol Methods* **128**: 183-188.
- Smal, M.A, Dziadek, M. and Cooney, S.J., Attard, M. and Baldo, B. A. 1990. Examination for platelet-activating factor production by preimplantation mouse embryos using a specific radioimmunoassay. *J Reprod Fertil*, **90**: 419-425.
- Smart, Y.C., Roberts, T.K., Clancy, R. and Cripps, A.W. 1981. Early pregnancy factor: its role in mammalian reproduction-research review. *Fertil Steril*, **35**:397-402.
- Snyder, F. 1990. Platelet-activating factor and related acetylated lipids as potent biologically active cellular mediators. *Am J Phys*, **259** C697-C708.
- Spinks, N.R. and O'Neill, C. 1988. Antagonists of embryo-derived platelet-activation factor prevent implantation of mouse embryos. *J Reprod Fertil*, **84**: 89-98.
- Sueoka, K., Dharmarajan, A.M., Miyazaki, T., Atlas, S.J. and Wallach, E.E. 1989. In vivo and in vitro determination of components of rabbit early pregnancy factors. *J Reprod Fertil*, **87**: 47-53.
- Tetta, C., Montrucchio, G., Alloati, G., Roffinello, C., Emanuello, G., Benedetto, C., Camussi, G. and Masobrio M. 1986. Platelet-activating factor contracts human myometrium in vitro. *Proc Soc Exp Biol Med* **183**: 376-381.
- Velasquez, L.A., Aguilera, J.G. and Croxatto, H.B. 1995. Possible role of platelet-activating factor in embryo signaling during oviductal transport in hamster. *Biol Reprod*, **52**:1302-1306
- Wallace, J.L. 1989. PAF as a mediator of gastrointestinal damage. In: Saito K, Hanahan DJ (eds.), *Platelet-activating factor and disease*. Tokyo: International Medical Publishers: 153-180.
- Wells, X.E. and O'Neill, C. 1994. Detection and preliminary characterization of two enzymes involved in biosynthesis of platelet-activating factor in mouse oocytes, zygotes and pre-implantation embryos: dethiothreitol-insensitive cytidinediphosphocholine:1-O-alkyl-2-acetyl-sn-glycerol cholinephosphotransferase and acetyl-coenzyme A:1-O-alkyl-2-lyso-sn-glycero-3- phosphocholine acetyltransferase. *J Reprod Fertil*, **101**:385-391.
- Wyhte, A. and Heap, R.B. 1983. Early pregnancy factor. *Nature*, **304**: 121-122
- Wykle, R.L., Malone, B. and Snyder, F. 1980. Enzymatic synthesis of 1-alkyl-2-acetyl-sn-glycero-3- phosphocholine, a hypotensive and platelet-aggregating lipid. *J Biol Chem*, **255**: 10256-10260.
- Yang, Y.Q., Kudolo, G.B., Harper, M.J.K. 1991. Platelet-activating factor (PAF) binding sites in the rabbit oviduct. *Arch Biol Med Exp* **24**: 275-279.
- Yang, Y.Q., Kudolo, G.B. and Harper, M.J.K. 1992. Binding of platelet-activating factor to oviductal membranes during early pregnancy in the rabbit. *J Lipid Mediators*, **5**: 77-96.
- Yasuda, K., Satouchi, K. and Saito, K. 1988. Platelet-activating factor in normal rat uterus. *Biochem Biophys Res Commun*, **138**:1231-1236

(Responsible Editor: Chai Ruihai)